

MINYAK ATSIRI DARI *Toona sinensis* DAN UJI AKTIVITAS INSEKTISIDA

Adlis Santoni¹, Hazli Nurdin¹, Yunazar Manjang¹, Sjamsul A. Achmad²

¹Jurusian Kimia FMIPA Universitas Andalas, Kampus Limau Manis, Padang

²Program Studi Kimia Institut Teknologi Bandung

email: adlis_1962@yahoo.com

ABSTRACT

Volatile components from leafs of *Toona sinensis* (Meliaceae) were isolated by distillation water-vapour method and analyzed using GC/MS. Forty-eight component were identified, and eight major components are Germacrene-D, Germacrene-B, α -Terpinene, α -Humulene, β -Caryophyllene, α -Elemene, Bicyclogermacrene and α -copaene. Insect activity of the essential oil against *Crosidolomia pavonana* were tested. The activity of the essential oil with concentration 25% (mortality 73.3%), concentration 50% (mortality 83.3%) and concentration 75% (mortality 90%).

Keywords : *Toona sinensis* (Meliaceae), essential oil, *Crosidolomia pavonana*, and mortality.

DAFTAR PUSTAKA

1. J. T. Arnason, S. Mackinnon, A. Durst, B. J. R. Philogene, Hasbun, P. Sanchez, L. Poveda, L. San Roman, M. B. Isman, C. Satasook, G. H. N Towers, P. Wiriachitra, and JLMC, Laughlin, Insecticides in tropical plants with non-neurotoxic modes of action, *Phytochemical Potential of Tropical Plants*, 107-151, (1993).
2. D. J. Mabberley, C. M. Pannel & A. M. Sing, *Flora Malesiana, Series I-Spermatophyta: Flowering plants*, Leiden University, 12: 1–20, 1995
3. SAS Institut, SAS/STAT User's Guide, North Carolina, SAS Institut Inc., Version 6, fourth ed., 2, 1990.
4. A. J. Prakash, Rao, Botanical pesticides in agriculture, CRC Press, New York, London, 1997, 461.
5. R. Nishida et al., *Journal of Chemical Ecology*, 26: 87, (2000).
6. T. Rostelien, A. K Borg-Karlson, The Plant Sesquiterpene Germacrene D Specifically Activates a major type of Antennal Receptor Neuron of the Tobacco Budworm Moth *Heliothis virescens*, *Chem. Sense*, 25: 141-148, (2000).
7. M. G. Chisholm, M. A. Wilson, et al., *Flavour and Fragrance Journal*. 18(2): 106-115, (2003).
- J. Lawrence, Porter, "Flavans and proanthocyanidins", In *The Flavanoids: Advances in Research Since 1986* (J.B. Harborne, Ed.), Chapman & Hall, London, 23–55, 1994.
- D. Ngokam, G. Massiot, J. M. Nuzillard and E. Tsamo, "(+)-7',7'-Dimethyl-5-hydroxy-2R,3S-Trans-Pubeschin From Entandrophragma cylindrum, *Phytochemistry*, 37: 529-531, (1994).
- D. E. Champagne, M. B. Isman, Towers GHN, *Insecticidal activity of phytochemicals and extracts of the Meliaceae*, Di dalam: Arnason JT, Philogene BJR, Morand P, editor. Insecticides of plant origin, Washington DC: ACS, 1989, 95-109.
- V. H. Kapadia et al., "Structure of mustakone and copaene". *Tetrahedron Letters*, 28: 1933, (1963).
- 1.

