

MULTIDRUG RESISTANCE (MDR) OF *V. Parahaemolyticus*

Marlina

Faculty of Pharmacy, University of Andalas
e-mail: marlina_adly@yahoo.com

ABSTRACT

A total of 97 *V. parahaemolyticus* isolate from Padang were examined for their resistance to 15 antibiotics. *V. parahaemolyticus* isolated behaved as resistant to sulfamethoxazole (100%), rifampin (95%) and tetracycline (75%) and sensitive to norfloxacin (96%). Ampicillin still sensitive for *V. parahaemolyticus* isolated from human stools. All of isolates were sensitive to namely chloramphenicol and fluoroquinolones (ciprofloxacin and norfloxacin agents). RAPD-PCR profiling with three primers (OPAR3, OPAR4 and OPAR8) produced four major clusters (R1, R2, R3 and R4), 7 minor clusters (I, II, III, IV, V, VI and VII) and three single isolates.

Keywords: *V. parahaemolyticus*, MDR, RAPD

REFERENCES

1. D. Ottaviani, I. Bacchicocchi, L. Masini, F. Leoni, A. Carraturo, M. Giammarioli, and G. Sbaraglia, Antimicrobial susceptibility of potentially halophilic vibrios isolated from seafood, *International Journal of Antimicrobial Agents* 18: 135-140, (2001).
2. A. Cespedes, and E. Larson, Knowledge, attitude and practices regarding antibiotic use among Latinos in the United States: Review and Recommendations, *American Journal of Infection Control* 34: 495-502, (2006).
3. M. Lesmana, D. Subekti, C.H. Simanjuntak, P. Tjaniadi, J. R. Campbell, and B. A. Ofoyo, *Vibrio parahaemolyticus* associated with cholera-like diarrhea among patients in North Jakarta, Indonesia, *Diagnostic Microbiology and Infectious Disease*, 39: 71-75, (2001).
4. S. Lu, B. Liu, B. Zhou, And R. E. Levin, Incidence and Enumeration of *Vibrio parahaemolyticus* in Shellfish from two retail Sources and the Genetic Diversity of isolates as Determined by RAPD-PCR Analysis, *Food Biotechnology*, 20: 193-209, (2006).
5. M. Nishibuchi, *Vibrio parahaemolyticus*. In *International handbook of foodborne pathogens*, ed. M.D. Milliots and J. W. Bier, United States: Marcel Dekker, Inc. P, 2004, 237-252.
6. L. Poirel, M. R. Martinez, H. Mammeri, A. Liard, and P. Nordmann, Origin of Plasmid-Mediated Quinolone Resistance Determinant QnrA, *Antimicrobial Agents and Chemotherapy*, 49: 3523-3525, (2005).
7. S. Radu, N. Elhadi, Z. Hassan, G. Rusul, S. Lihan, N. Fifadara, Yuherman and E. Purwati, Characterization of *Vibrio vulnificus* isolated from cockles (*Anadara granosa*): antimicrobial resistance, plasmid profiles and random amplification of polymorphic DNA analysis, *FEMS Microbiology Letters*, 165: 139-143, (1998).
8. S. Radu, N. Ahmad, F. H. Ling, and A. Reezal, Prevalence and resistance to antibiotics for *Aeromonas* species from retail fish in Malaysia, *International Journal Food Microbiology*, 81: 261-266, (2003).

9. B. Sarkar, N. R. Chowdhury, G. B. Nair, M. Nishibuchi, S. Yamasaki, Y. Takeda, S. K. Gupta, S. K. Bhattacharya, and Ramamurthy, Molecular characterization of *Vibrio parahaemolyticus* of similar serovars isolated from sewage and clinical cases of diarrhea in Calcutta, India, *World Journal of Microbiology and Biotechnology*, 19: 771-776, (2003).
10. S. Schwarz, and E. Chaslus-Dancla, Use of antimicrobials in veterinary medicine and mechanisms of resistance, *Veterinary Residue*, 32: 201–225, (2001).
11. H. Sörum, and T.M. L'Abée-Lund, Antibiotic resistance in food-related bacteria – a result of interfering with the global web of bacterial genetics, *International Journal of Food Microbiology*, 78: 43–56, (2002).
12. P. Tjaniadi, M. Lesmana, D. Subekti, N. Machpud, S. Komalarini, W. Santoso, C. H. Simanjuntak, N. Punjabi, J. R. Campbell, W. K. Alexander, H. J. Beecham, A. L. Corwin, and B. A. Oyofo, Antimicrobial Resistance of Bacterial Pathogens Associated with Diarrheal Patients in Indonesia, *American Journal of Tropical Medicine and Hygiene*, 68: 666-670, (2003).
13. X. Zhao, and D. Drlica, Restricting the Selection of Antibiotic-Resistant Mutants: A General Strategy Derived from Fluoroquinolone Studies, *Clinical Infectious Diseases*, 33: S147-S156, (2001).